

Self-Dual metrics on Non-simply connected 4-Manifolds

Hülya Argüz

Mustafa Kalafat

Yıldırım Ozan

July 18, 2012

Abstract

We construct self-dual(SD) but not locally conformally flat(LCF) metrics on families of non-simply connected 4-manifolds with small signature. We construct various sequences with bounded or unbounded Betti numbers and Euler characteristic. These metrics have negative scalar curvature. As an application, this addresses the Remark 4.79 of [Bes].

1 Introduction

Self-dual manifolds were introduced by Penrose in [Pen], and they are put on a firm mathematical foundation by Atiyah, Hitchin and Singer in [AHS]. After the explicit constructions of Poon, LeBrun, Joyce, and the connected sum theorem of Donaldson and Friedman, Taubes proved that every 4-manifold admits an anti-self-dual metric after taking connected sum with $k\overline{\mathbb{CP}}_2$, where k is sufficiently large [Tau]. Although this is a very useful theorem, the anti-self-dual metric here is not explicit. The minimal number for k is called the *Taubes Invariant*, which is unknown for most 4-manifolds. The number k in Taubes' work is very large. There are many explicit constructions of self-dual metrics on simply connected manifolds, however there are very few examples in the non-simply connected case which are not locally conformally flat. The self-dual metrics on the blow ups of the connected sums of $S^3 \times S^1$ [LeH, Kim], and on the blow ups of $S^2 \times \Sigma_g$ for $g \geq 1$ [LeR, KLP] are the only explicitly known examples. Here we give new examples of self-dual metrics on closed non-simply connected 4-manifolds, and show that many new topological types can be realized.

The idea is first to construct LCF manifolds using the techniques introduced in [AK]. See Figure 1 for the construction of the manifolds. The manifolds in [AK] does not satisfy the hypothesis to produce self-dual metrics with minimal number of blow up. So one needs to make modifications and produce a new family. One needs to disconnect the boundary of the related 3-manifold. So that J.Kim's theorems are applicable and as a consequence these manifolds admit hyperbolic ansatz self-dual metrics of LeBrun's. We obtain the following results.

Theorem 1.1. *The manifolds $M_{g,n}^5 \# l\mathbb{CP}_2$ admits self-dual metric of negative scalar curvature for all $l \geq 2$.*

Corollary 1.2. *There are new infinite families of closed, non-simply connected, self-dual 4-manifolds, with betti number growth as follows.*

1. $b_1 \rightarrow \infty$, b_2 bounded, and $\chi \rightarrow -\infty$,
2. $b_1 \rightarrow \infty$, $b_2 \rightarrow \infty$, and $\chi \rightarrow -\infty$,
3. $b_1 \rightarrow \infty$, $b_2 \rightarrow \infty$, and χ bounded,
4. b_1 bounded, $b_2 \rightarrow \infty$, and $\chi \rightarrow \infty$,
5. $b_1 \rightarrow \infty$, $b_2 \rightarrow \infty$, and $\chi \rightarrow \infty$.

These manifolds have strictly negative scalar curvature.

We also prove the following theorem using Kim's result. Note that this can not be obtained from Taubes' theorem since it does not say anything about the scalar curvature. See [AK] for the constructions of the manifolds in this theorem.

Theorem 1.3. *The manifolds $M_g^1 \# l\mathbb{CP}_2$, $M_{g,n}^2 \# l\mathbb{CP}_2$, $M_1^4 \# l\mathbb{CP}_2$ admits self-dual metric of negative scalar curvature for all sufficiently large l .*

Since their signature is nonzero, the above manifolds do not admit any locally conformally flat(LCF) metrics. In the Remark 4.79, [Bes] asks for examples of compact, self-dual (half-conformally flat) manifolds which are not conformally Einstein. The family of metrics in the parts 1 and 2 of the Corollary 1.2 has negative Euler characteristic. This violates the Hitchin-Thorpe inequality, see [B] or the more recent [T, H]. If there is an Einstein or conformally Einstein metric on these compact 4-manifolds then the Euler characteristic has to be non-negative by Hitchin-Thorpe. So that these metrics are instances in the remark which are explicit, non LCF and of negative scalar curvature.

Acknowledgements. We would like to thank to Claude LeBrun for directing us to the field, Jongsu Kim for his encouragement. The figure is constructed by the IPE software of Otfried Cheong.

2 Background and Proofs

To be able to construct the self-dual metrics, we first construct locally conformally flat(LCF) metrics using Braam's conformal compactification procedure in [Br]. One starts with a hyperbolic 3-manifold N with boundary which is obtained from the hyperbolic space by taking the quotient with a cusp-free geometrically finite Kleinian group. Then one spins around the boundary to get a closed Riemannian 4-manifold X ,

$$X := \frac{N \times S^1}{\{b \times S^1 \mid b \in \partial N\}}.$$

This is to cross with a circle and then contracting the circles that multiplied with a boundary point. This process coupled with the magnetic monopoles of LeBrun [LeEx] called the hyperbolic ansatz yields self-dual metrics on the blow ups. The following theorem of J.Kim tells the precise conditions to be able to carry out the hyperbolic ansatz.

Theorem 2.1 ([Kim]). *If $N = \mathcal{H}^3/\Gamma$ is a noncompact hyperbolic 3-manifold obtained from a cusp-free geometrically finite Kleinian group Γ and $H_2(\bar{N}; \mathbb{R})/H_2(\partial N; \mathbb{R})$ is at most 1-dimensional, then there exist self-dual metrics on $X \sharp l\mathbb{CP}_2$ for all sufficiently large natural number l where X is obtained from N by spinning around its boundary. The sign of the conformal class of these metrics are identical to the sign of $1-d(\Lambda(\Gamma))$.*

If moreover $H_2(\bar{N}; \mathbb{R}) = H_2(\partial N; \mathbb{R})$ we can take l to be bigger than or equal to the number of connected components of ∂N .

To check the homological condition appearing in the hypothesis of the theorem, we need to use the following isomorphism [Br] where the coefficients are integers.

$$H_2(\bar{N}) \oplus H_1(\bar{N}, \partial \bar{N}) \xrightarrow{\sim} H_2(X). \quad (1)$$

To establish this isomorphism one needs a little bit of work, details of which is not given in [Br] so we will prove it here. Rather than shrinking the boundary circles, one can attach 2-discs to fill out those circles, called the capping [AK] operation. So one can work with the following decomposition of the 4-manifold.

$$X = \bar{N} \times S^1 \cup \partial \bar{N} \times D^2 \quad \text{where} \quad \bar{N} \times S^1 \cap \partial \bar{N} \times D^2 = \partial \bar{N} \times S^1.$$

If we write down the Mayer-Vietoris exact sequence for this pair we get the following piece.

$$\cdots \rightarrow H_2(\partial N \times S^1) \xrightarrow{A} H_2(N \times S^1) \oplus H_2(\partial N) \xrightarrow{B} H_2(X) \xrightarrow{C} H_1(\partial N \times S^1) \xrightarrow{D} H_1(N \times S^1) \oplus H_1(\partial N) \rightarrow \cdots$$

To analyze this sequence we need the maps involved in the relative exact sequence.

$$\cdots \rightarrow H_1(\partial N) \xrightarrow{i_*^1} H_1(N) \xrightarrow{j_*^1} H_1(N, \partial N) \xrightarrow{\partial_*^1} H_0(\partial N) \xrightarrow{i_*^0} H_0(N) \rightarrow 0.$$

From this point on, under the assumption that ∂N has one or two connected components, one gets an easier proof. Nonetheless we will prove the general assertion. We make up the following exact diagram.

$$\begin{array}{ccccc}
& & H_2(X) & \xrightarrow{C} & \text{Ker } i_*^0 \\
& \nearrow B & & & \searrow \\
H_2(\partial N) \oplus H_1(\partial N) & \xrightarrow{A} & H_2(N) \oplus H_1(N) \oplus H_2(\partial N) & & 0 \\
& \searrow \varphi & & & \nearrow \\
& & H_2(N) \oplus \text{Im } j_*^1 & &
\end{array}$$

The domain and range of the map A is decomposed into basic components according to the Künneth formula, and is given by $A(a, b) = (i_*a, i_*b, -a)$. Using these conventions we define the new map φ by $\varphi(x, y, z) = (x + i_*z, j_*^1y)$. One can easily check the exactness by chasing through the relative exact sequence, where the map j_*^1 is also involved. So that $\text{Ker } \varphi = \text{Im } A$ and φ is surjective. The upper part is taken from the Mayer-Vietoris sequence above, where the map D is defined by $D = I \oplus -I$, and after decomposing into components with respect to Künneth, one can make the identification $\text{Ker } D = \text{Ker } i_*^0$. Since $\text{Coker } A \approx \text{Im } \varphi$, the short exact sequence with nucleus $H_2(X)$ in Mayer-Vietoris reads

$$0 \longrightarrow H_2(N) \oplus \text{Im } j_*^1 \xrightarrow{\bar{B}} H_2(X) \xrightarrow{C} \text{Ker } i_*^0 \longrightarrow 0.$$

On the other hand, since the $\text{Coker } i_*^1 \approx \text{Im } j_*^1$, pulling out the short exact sequence with nucleus $H_1(N, \partial N)$ from the relative sequence reads,

$$0 \longrightarrow \text{Im } j_*^1 \xrightarrow{\text{inc}} H_1(N, \partial N) \xrightarrow{\partial_*^1} \text{Ker } i_*^0 \longrightarrow 0.$$

All the maps are natural. Now that $H_0(\partial N)$ is always free so that these exact sequences split, and combining the two yields the isomorphism (1).

Next using the Lefschetz duality $H_1(N, \partial N) \approx H^2(N)$ and universal coefficients theorem, after cancelling the Ext terms the isomorphism becomes,

$$\text{Hom}(H_2(N), \mathbb{Z}) \oplus H_2(N) \approx \text{Hom}(H_2(X), \mathbb{Z}).$$

Writing according to the free and torsion pieces it looks like

$$F_2N \oplus F_2N \oplus T_2N \approx F_2X.$$

So that comparing both sides we obtain $T_2N = 0$ and $2F_2N \approx F_2X$.

Now the panelled web 4-manifolds M_g^1 , $M_{g,n}^2$ and M_1^4 of [AK] has $b_2 = 2$. By above this gives $b_2(N) = 1$ which satisfies the first hypothesis of the Theorem 2.1. However, they do not satisfy the second homological hypothesis. Next we construct a new set of manifolds to handle this situation. We modify the sequence $M_{g,n}^2$. To satisfy the

second hypothesis, the boundary surfaces should generate the second homology of the 3-manifold N . The examples in [AK] has all connected boundaries, and the boundary bounds the 3-manifolds and homologous to zero, hence the map

$$i_* : H_2(\partial N, \mathbb{Z}) \rightarrow H_2(N, \mathbb{Z})$$

is zero. In order to get a nontrivial image, we should disconnect the boundary. So whenever identifying the boundary cylinders in the construction of $M_{g,n}^2$'s we identify in the parallel way this time to get two distinct boundary components. This way we obtain the sequence of manifolds $M_{g,n}^5$ which is shown concretely in Figure 1. The




Figure 1: The LCF manifolds $M_{g,n}^5$.

following theorem ensures that the boundary surfaces generate $H_2(N, \mathbb{Z})$.

Theorem 2.2. *Let N be a compact orientable 3-manifold with two boundary components and $b_2 = 1$. Then each of the boundary surfaces is a generator of $H_2(N, \mathbb{Z})$.*

Proof. Take two copies of the manifold N_1 and N_2 so that the second copy is reversely oriented. Then identify them along their boundaries to form the compact 3-manifold $M = N_1 \cup_{\partial N} N_2$ without boundary, which we call *the double* of N . If we look at the Mayer-Vietoris exact sequence in this case, we see the following piece.

$$\begin{array}{ccccccc} 0 \longrightarrow H_3(M) \longrightarrow H_2(\partial N) & \xrightarrow{i_*^1 \oplus -i_*^2} & H_2(N_1) \oplus H_2(N_2) & \xrightarrow{j_*^1 + j_*^2} & H_2(M) \longrightarrow \cdots \\ & \wr & & & \\ & \mathbb{Z} \oplus \mathbb{Z} & & & \mathbb{Z} \oplus \mathbb{Z} \end{array}$$

The second isomorphism here can be checked through the universal coefficients theorem that there is no torsion. Assuming that the maps acts through the canonical isomorphisms, the image of the generators are,

$$i_*^1 \oplus -i_*^2 : (1, 0) \mapsto (k, k), \quad (0, 1) \mapsto (l, l)$$

for some $k, l \in \mathbb{Z}$. Since ∂N bounds both N_1 and N_2 , the image of $(1, 0) + (0, 1)$ under i_*^1 and i_*^2 is zero implying $l = k$. Since by compactness $H_2(M)$ is torsion free, the isomorphism following from exactness

$$\mathbb{Z} \oplus \mathbb{Z} / \langle k(1, 1) \rangle \approx H_2(M)$$

forces $k = \pm 1$, hence a generator in both components. Furthermore, we have $H_2(M) = \mathbb{Z}$. $\square$

Since both of the homological conditions of [Kim] are satisfied, Theorem 1.1 follows.

Next we compute the topological invariants. The fundamental group,

$$\langle a_1 \cdots b_g, c_1 \cdots c_n, d_1 \cdots d_n, e_1 \cdots e_n \mid a_1^{-1} b_1^{-1} a_1 b_1 \cdots a_g^{-1} b_g^{-1} a_g b_g c_1 \cdots c_n d_n^{-1} \cdots d_1^{-1}, e_i d_i e_i^{-1} c_i \rangle$$

gives relations $d_i = c_i^{-1}$ for $i = 1 \cdots n$ and $(c_1 \cdots c_n)^2 = 1$, after abelianization. Introducing a new variable $\bar{c} := c_1 \cdots c_n$ and dropping the redundant $c_n = c_{n-1}^{-1} \cdots c_1^{-1} \bar{c}$ yields

$$H_1(M_{g,n}^5; \mathbb{Z}) = \langle a_1 \cdots b_g, c_1 \cdots c_{n-1}, \bar{c}, e_1 \cdots e_n \mid \bar{c}^2 \rangle_+ \approx \mathbb{Z}^{2g+2n-1} \oplus \mathbb{Z}_2.$$

Counting the handles $\chi = 4 - 4g - 4n$ and $b_2 = 0$, then $H_2(M_{g,n}^5; \mathbb{Z}) = \mathbb{Z}_2$. After connecting summing with l copies of $\mathbb{CP}_2$, we get for the manifolds $M_{g,n}^5 \# l\mathbb{CP}_2$

$$b_1 = 2g + 2n - 1, \quad b_2 = l, \quad \chi = 4 - 4g - 4n + l.$$

Now, the sequences in the Corollary 1.2 can be obtained by letting $g, n \rightarrow \infty$, and then taking $l = g + 2$ or $l = 4(g + n)$ in addition. Letting $l \rightarrow \infty$, and then taking $l = 5(g + n)$ in addition. Alternatively, taking $l = g + n$, one can again produce other sequences of the second type.

UNIVERSITÄT HAMBURG, BUNDESSTRASSE 55, 20146, GERMANY
E-mail address: huelya.arguez@math.uni-hamburg.de

TUNCELİ ÜNİVERSİTESİ, TURKIA
E-mail address: kalafg@gmail.com

ORTA DOĞU TEKNİK ÜNİVERSİTESİ, 06800, ANKARA, TURKIA
E-mail address: ozan@metu.edu.tr

References

- [AK] S. AKBULUT, M. KALAFAT, *Some new locally conformally flat 4-manifolds*. Preprint available at Arxiv math.DG/0807.0837.
- [AHS] M. F. ATIYAH, N. J. HITCHIN, AND I. M. SINGER, *Self-duality in four-dimensional Riemannian geometry*. Proc. Roy. Soc. London Ser. A, 362 (1978), pp. 425–461.
- [B] M. BERGER, *Sur quelques variétés d'Einstein compactes. (French)*. Ann. Mat. Pura Appl. (4) 53. 1961. 89–95.
- [Bes] A. L. BESSE, *Einstein Manifolds*, Reprint of the 1987 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2008. xii+516 pp. ISBN: 978-3-540-74120-6.
- [BisJon] C. BISHOP, P. JONES, *Hausdorff dimension and Kleinian groups*, Acta Math. 179 (1997), no. 1, 1-39.
- [Br] P. J. BRAAM, *A Kaluza-Klein approach to hyperbolic three-manifolds*, Enseign. Math. (2) 34 (1988), no. 3-4, 275–311.
- [CaMiTa] R. CANARY, Y. MINSKY, E. TAYLOR, *Spectral theory, Hausdorff dimension and the topology of hyperbolic 3-manifolds*, J. Geom. Anal. 9 (1999), no. 1, 17-40.
- [GS] R. E. GOMPF AND A. I. STIPSICZ, *4-Manifolds and Kirby Calculus*, American Mathematical Society, Providence, RI, 1999.
- [Ha] A. HATCHER, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. xii+544 pp. ISBN: 0-521-79160-X.
- [H] N. J. HITCHIN, *Compact four-dimensional Einstein manifolds*. J. Differential Geometry 9. (1974), 435-441.
- [Kim] JONGSU KIM, *On the scalar curvature of self-dual manifolds*, Math. Ann. 297 (1993), no. 2, 235–251.
- [KLP] J. KIM, C. LEBRUN, M. PONTECORVO, *Scalar-flat Kähler surfaces of all genera*, J. Reine Angew. Math. 486 (1997), 69–95.
- [LeH] C. LEBRUN, *Anti-self-dual Hermitian metrics on blown-up Hopf surfaces*, Math. Ann. 289 (1991), no. 3, 383–392.
- [LeOM] C. LEBRUN, *Curvature Functionals, Optimal Metrics, and the Differential Topology of 4-Manifolds*, in Different Faces of Geometry, Donaldson, Eliashberg, and Gromov, editors, Kluwer Academic/Plenum, 2004.

- [LeEx] C. LEBRUN, *Explicit self-dual metrics on $CP_2\# \cdots\# CP_2$* . J. Differential Geom. 34 (1991), no. 1, 223–253.
- [LeR] C. LEBRUN, *Scalar-flat Kähler metrics on blown-up ruled surfaces*, J. Reine Angew. Math. 420 (1991), 161–177.
- [MaPG] B. MASKIT, *Panelled web groups*, Kleinian groups and related topics (Oaxtepec, 1981), pp. 79–108, Lecture Notes in Math., 971, Springer, Berlin-New York, 1983.
- [Pen] R. PENROSE, *Nonlinear gravitons and curved twistor theory*, The riddle of gravitation—on the occasion of the 60th birthday of Peter G. Bergmann (Proc. Conf., Syracuse Univ., Syracuse, N. Y., 1975). General Relativity and Gravitation 7 (1976), no. 1, 31–52.
- [Tau] C.H. TAUBES, *The existence of anti-self-dual conformal structures*, J. Differential Geom. 36 (1992), no. 1, 163–253.
- [T] J. A. THORPE, *Some remarks on the Gauss-Bonnet integral*. J. Math. Mech. 18. 1969. 779–786.